arXiv:1607.08850v1 [math.CO] 29 Jul 2016

Bounding the distance among longest paths in a connected graph

Jan Ekstein* Shinya Fujita† Adam Kabela* Jakub Teska*

August 1, 2016

Abstract

It is easy to see that in a connected graph any 2 longest paths have a vertex in common. For $k \geq 7$, Skupień in [7] obtained a connected graph in which some k longest paths have no common vertex, but every $k-1$ longest paths have a common vertex. It is not known whether every 3 longest paths in a connected graph have a common vertex and similarly for $4, 5$, and 6 longest path. In [5] the authors give an upper bound on distance among 3 longest paths in a connected graph. In this paper we give a similar upper bound on distance between 4 longest paths and also for k longest paths, in general.

1 Introduction

In 1966 Gallai in [4] asked whether all longest paths in a connected graph have a vertex in common. Couple of years later, several counterexamples were found, see [9], [10], and [11]. In 1976 Thomassen in [8] showed that there exist infinitely many counterexamples to Gallai's question.

On the other hand, if we restrict to a special class of graphs, the answer to Gallai's question may become positive. For example in a tree, all longest paths must have a vertex in common. Klavžar and Petkovšek in [6] proved that it is also true for split graphs and cacti and Balister et al. in [2] proved it for the class of circular arc graphs.

Another approach to Gallai's question is to ask, what happens if we consider a fixed number of longest paths. It is easy to see that every 2 longest paths in a connected graph have a common vertex. For 3 longest paths, the question remains open. This has been originally asked by Zamfirescu in [12].

*Department of Mathematics, Institute for Theoretical Computer Science, and European Centre of Excellence NTIS - New Technologies for the Information Society, Faculty of Applied Sciences, University of West Bohemia, Pilsen, Technická 8, 306 14 Plzeň, Czech Republic, e-mail: {ekstein,kabela,teska}@kma.zcu.cz.

†International College of Arts and Sciences, Yokohama City University, 22-2 Seto, Kanazawa-ku, Yokohama 236-0027, Japan, e-mail: shinya.fujita.ph.d@gmail.com.

Conjecture 1. [12] *For every connected graph, any* 3 *of its longest paths have a common vertex.*

There are few results dealing with this conjecture. Axenovich in [1] proved that it is true for connected outerplanar graphs and de Rezende et al. in [3] showed that Conjecture 1 is true for connected graphs in which all nontrivial blocks are hamiltonian.

For $k \geq 7$, Skupień in [7] obtained a connected graph in which some k longest paths have no common vertex, but every $k-1$ longest paths have a common vertex. Regarding this, it is still valid to ask wheter not only 3 but also $4, 5$, and 6 longest path in a connected graph have a common vertex.

In [5] the authors introduced a parameter to measure the distance among the longest paths in a connected graph and proved an upper bound of this parameter for 3 longest paths. To state their result we give some definitions first.

Let G be a connected graph. Let $\ell(G)$ be the length of any longest path in G and $\mathcal{L}(G) = \{P \mid P \text{ is a path in } G \text{ with } |V(P)| = \ell(G)+1\}$ be a set of longest paths of G. For $x, y \in V(G)$, let $d_G(x,y)$ be a distance between x and y in G. For a vertex $x \in V(G)$ and a subset $U \subseteq V(G)$, let $d_G(x,U) = min\{d_G(x,y) |\ y \in U\}$. For $\mathcal{P} \subseteq \mathcal{L}(G)$ we call *a path-distance-function* $f(G,\mathcal{P}) = min\{\sum_{P\in\mathcal{P}} d_G(v,V(P)) \mid v \in V(G)\}$.

For a class of graphs $\mathcal{G}$ and an integer k, we introduce *a path-distance-ratio* $d_k(\mathcal{G}) = max\frac{f(G,\mathcal{P})}{|V(G)|}$, where the maximum is taken overall graphs of $\mathcal{G}$ and their sets of longest paths $\mathcal{P} \subseteq \mathcal{L}(G)$ with $|\mathcal{P}| = k$.

Let $\mathcal{G}_c$ be a class of connected graphs. The question whether for every connected graph any 3 longest paths have a vertex in common translates into a question whether $d_3(\mathcal{G}_c) = 0$. On the other hand, Skupień in [7] constructed a graph on 17 vertices, in which there are 7 longest paths without a common vertex, this graph implies that $d_7(\mathcal{G}_c) \geq \frac{1}{17}$.

Now we can state the result by Fujita et al. from [5].

Theorem 2. [5] *Let $\mathcal{G}_c$ be a class of connected graphs. Then $d_3(\mathcal{G}_c) \leq \frac{1}{17}$.*

In this paper we prove similar results for 4 longest path and also for k longest paths, in general.

Theorem 3. *Let $\mathcal{G}_c$ be a class of connected graphs. Then $d_4(\mathcal{G}_c) \leq \frac{3}{16}$.*

By picking any vertex of a connected graph G, we see that $d_k(\mathcal{G}_c)$ can be bounded by k. We show that it can be improved as roughly $\frac{k}{6}$.

Theorem 4. *Let $\mathcal{G}_c$ be a class of connected graphs and let $k \geq 3$ be an integer. Then $d_k(\mathcal{G}_c) \leq \frac{k^3-4k^2+5k-2}{6k^2-8k}$.*

2 Proofs

In our proofs, we adapt ideas of [5]. We start by giving several technical definitions.

Let G be a connected graph. Let P be a path in G and U and V be two sets of vertices of G, a $U-V$ path on P is a path on P with end-vertices u and v such that $u \in U$ and $v \in V$. We call *int-vertices* all vertices of P except u and v. Let uPv denote the $\{u\}-\{v\}$ path on P. Futhermore, let $\breve{u}Pv = uPv - u$, $uP\breve{v} = uPv - v$ and $\breve{u}P\breve{u} = uPv - \{u, v\}$. For a set $\mathcal{P} = \{P, P_1, P_2, ..., P_{k-1}\} \subseteq \mathcal{L}(G)$ and $i \neq j \in \{1, 2, ..., k-1\}$, a $V(P_i) - V(P_j)$ path Q on P is *good* if $V(Q) \cap V(P_m) \neq \emptyset$ for every $m = 1, 2, ..., k-1$ and neither P_i nor P_j contain an int-vertex of Q. Let $t_{\mathcal{P}}(P)$ be the number of all good paths of P and $t'_{\mathcal{P}}(P)$ be the maximum number of all non-intersecting (no edge in common) good paths on P. By Proposition 3 in [5] (every 2 longest paths intersect) we have that $t_{\mathcal{P}}(P) \geq t'_{\mathcal{P}}(P) \geq 1$ for every $P \in \mathcal{P}$. For a path $P \in \mathcal{P}$, let $X^i_{\mathcal{P}}(P)$ denote the set of all vertices of P which are exactly on i paths from $\mathcal{P}$. Let $n_i = |\bigcup_{P \in \mathcal{P}} X^i_{\mathcal{P}}(P)|$.

Lemma 5. *Let G be a connected graph of order n and $\mathcal{P} \subseteq \mathcal{L}(G)$ with $|\mathcal{P}| = k \geq 3$. If $f(G, \mathcal{P}) > 0$, then*

$$n \geq \frac{k \cdot \ell(G) + k + (k-2)n_1 + (k-3)n_2 + ... + n_{k-2}}{k-1}.$$

Proof. Clearly $n \geq n_1 + n_2 + ... + n_{k-1} + n_k$, where $n_k = 0$, and $n \geq k(\ell(G)+1) - n_2 - 2n_3 - ... - (k-3)n_{k-2} - (k-2)n_{k-1}$. Hence $n \geq k \cdot \ell(G) + k - n_2 - 2n_3 - ... - (k-3)n_{k-2} - (k-2)(n - n_1 - n_2 - ... - n_{k-2})$ and the result follows. □

Lemma 6. *Let G be a connected graph and $\mathcal{P} \subseteq \mathcal{L}(G)$ with $|\mathcal{P}| = k$. If there exists a path $P \in \mathcal{P}$ with $t'_{\mathcal{P}}(P) = 1$, then $f(G, \mathcal{P}) = 0$.*

Proof. To the contrary, we suppose there is a path $P = v_1 v_2 ... v_{\ell(G)+1}$ with $t'_{\mathcal{P}}(P) = 1$ and $f(G, \mathcal{P}) > 0$. By $f(G, \mathcal{P}) > 0$, every good path on P contains an edge. We consider the 'left-most' good path Q on P; more formally, we consider the good path $Q = v_i v_{i+1} ... v_j$ such that there is no good path on P containing a vertex v_k with $k < i$. Let $\mathcal{P}_j$ denote the set of paths of $\mathcal{P}$ which contain v_j. By the choice of Q, some path of $\mathcal{P}_j$ contains no vertex v_k with $k < j$, and thus the length of $v_1 v_2 ... v_j$ is at most $\frac{1}{2}\ell(G)$. Similarly, we consider the 'right-most' good path $Q' = v_{i'} v_{i'+1} ... v_{j'}$ and we see that the length of $v_{i'} v_{i'+1} ... v_{\ell(G)+1}$ is at most $\frac{1}{2}\ell(G)$. By the assumption $t'_{\mathcal{P}}(P) = 1$, the paths Q and Q' have an edge in common, so $j > i'$, hence the length of P is shorter than $\ell(G)$, a contradiction. □

Lemma 7. *Let G be a connected graph and $\mathcal{P} \subseteq \mathcal{L}(G)$ with $|\mathcal{P}| = k \geq 3$. Let $P \in \mathcal{P}$ and let Q be a good path on $\mathcal{P}$. Then the following two statements hold:*

(i) $f(G, \mathcal{P}) \leq \frac{|V(Q)|-1}{2}(k-1)$;

(ii) $|X^1_{\mathcal{P}}(P) \cup X^2_{\mathcal{P}}(P) \cup ... \cup X^{k-2}_{\mathcal{P}}(P)| \geq t'_{\mathcal{P}}(P)(\frac{2}{k-1} f(G, \mathcal{P}) - 1)$.

Proof. Note that if $f(G, \mathcal{P}) = 0$, then the statement holds. Suppose $f(G, \mathcal{P}) \geq 1$. In particular, every good path on $\mathcal{P}$ contains at least two vertices. Let $x \in V(Q)$ such that $\sum_{P' \in \mathcal{P}} d_G(x, P') \leq \sum_{P' \in \mathcal{P}} d_G(y, P')$ for every $y \in V(Q)$. Then

$$f(G, \mathcal{P}) \leq \sum_{P' \in \mathcal{P}} d_G(x, P') \leq \frac{|V(Q)| - 1}{2}(k-1).$$

For any path P of $\mathcal{P}$ and any good path Q' on P, no int-vertex of Q' is in $X^{k-1}_{\mathcal{P}}(P)$, therefore $|V(Q') \cap (|X^1_{\mathcal{P}}(P) \cup X^2_{\mathcal{P}}(P) \cup ... \cup X^{k-2}_{\mathcal{P}}(P))| \geq |V(Q')| - 2 \geq \frac{2}{k-1} f(G,\mathcal{P}) - 1$. Let $\mathcal{Q}$ be a maximum set of non-intersecting good paths on P. By definition, $t'_{\mathcal{P}}(P) = |\mathcal{Q}|$, and we have

$$|X^1_{\mathcal{P}}(P) \cup X^2_{\mathcal{P}}(P) \cup ... \cup X^{k-2}_{\mathcal{P}}(P)| \geq |\cup_{Q \in \mathcal{Q}} (V(Q) \cap (X^1_{\mathcal{P}}(P) \cup X^2_{\mathcal{P}}(P) \cup ... \cup X^{k-2}_{\mathcal{P}}(P)))| \geq$$

$$\geq \sum_{Q \in \mathcal{Q}} (|V(Q)| - 2) \geq t'_{\mathcal{P}}(P) \left(\frac{2}{k-1} f(G,\mathcal{P}) - 1 \right).$$

□

Corollary 8. *Let G be a connected graph and $\mathcal{P} \subseteq \mathcal{L}(G)$ with $|\mathcal{P}| = 4$. Let $\mathcal{P} = \{P, P_1, P_2, P_3\}$ and let Q be a good path on $\mathcal{P}$. Then the following two statements hold:*

(i) $f(G,\mathcal{P}) \leq |V(Q)| - 1$;

(ii) $|X^1_{\mathcal{P}}(P) \cup X^2_{\mathcal{P}}(P)| \geq t'_{\mathcal{P}}(P)(f(G,\mathcal{P}) - 1)$.

Proof. The proof is the same as the proof of Lemma 7 with respect to the following. Let u, v be end-vertices of Q. Assume that Q is a $V(P_1) - V(P_2)$ path on P (otherwise we renumber the paths) and we consider a vertex $x \in V(Q) \cap V(P_3)$. Then

$$f(G,\mathcal{P}) \leq \sum_{P \in \mathcal{P}} d_G(x,P) = d_G(x,P_1) + d_G(x,P_2) \leq d_G(u,v) \leq |V(Q)| - 1.$$

Then we use Corollary 8(i) instead of Lemma 7(i) and the result follows. □

Proof of Theorem 4. Suppose that $f(G,\mathcal{P}) \geq 1$. Hence $t'_{\mathcal{P}}(P) \geq 2$ by Lemma 6. Let $P \in \mathcal{P}$ be a path minimizing $|X^1_{\mathcal{P}}(P) \cup X^2_{\mathcal{P}}(P) \cup ... \cup X^{k-2}_{\mathcal{P}}(P)|$. Let $\mathcal{P} - \{P\} = \{P_1, P_2, ..., P_{k-1}\}$ and u_i, v_i be the end-vertices of P_i for $i \in \{1, 2, ..., k-1\}$. Assume that Q is a good $V(P_1) - V(P_2)$ path on P with end-vertices u, v (otherwise we renumber paths $P_1, P_2, ..., P_{k-1}$). Let R be the shortest $\{u\} - V(P_2)$ path on P_1 and $x \in V(R) \cap V(P_2)$. We may assume that $|V(u_2P_2v)| \leq |V(u_2P_2x)|$ (see Figure 1).

We have $|V(R)| \geq 2$ from $f(G,\mathcal{P}) \geq 1$ and $|V(Q)| \geq \frac{2f(G,\mathcal{P})}{k-1} + 1$ from Lemma 7(i). Since $vQ\check{u}$ contains no vertex of $V(P_1)$, $vQuRx$ is a path in G. Futhermore, since $\check{v}QuP_1\check{x}$ contains no vertex of $V(P_2)$, $S_1 = v_2P_2vQuR\check{x}$, $S_2 = u_2P_2vQuRxP_2v_2$, and $S_3 = u_2P_2xRuQ\check{v}$ are paths in G (see Figure 2).

By comparing the lengths of P_2 and S_1 and using Lemma 7(i) and $|V(R)| \geq 2$, we have

$$|V(u_2P_2v)| - 1 \geq |V(Q)| - 1 + |V(R)| - 2 \geq |V(Q)| - 1 \geq \frac{2f(G,\mathcal{P})}{k-1}.$$

Similarly for P_2 and S_2, we have

$$|V(vP_2x)| - 1 \geq |V(Q)| - 1 + |V(R)| - 1 \geq |V(Q)| \geq \frac{2f(G,\mathcal{P})}{k-1} + 1.$$



Figure 1: A good $V(P_1) - V(P_2)$ path Q and path R

Also for P_2 and S_3, we have

$$|V(xP_2v_2)| - 1 \geq |V(Q)| - 1 + |V(R)| - 2 \geq |V(Q)| - 1 \geq \frac{2f(G,\mathcal{P})}{k-1}.$$

Therefore all together we have

$$\ell(G) = |V(P_2)| - 1 = |V(u_2P_2v)| - 1 + |V(vP_2x)| - 1 + |V(xP_2v_2)| - 1 \geq$$

$$\geq \frac{2f(G,\mathcal{P})}{k-1} + \frac{2f(G,\mathcal{P})}{k-1} + 1 + \frac{2f(G,\mathcal{P})}{k-1} = \frac{6f(G,\mathcal{P})}{k-1} + 1. \qquad (*)$$

Clearly $n_i = \frac{1}{i}\sum_{P'\in\mathcal{P}} X^i_{\mathcal{P}}(P')$. By the choice of P and $t'_{\mathcal{P}}(P') \geq 2$ for every $P' \in \mathcal{P}$ together with $(*)$, Lemma 5, and Lemma 7 we have

$$n \geq \frac{k\cdot\ell(G) + k + (k-2)\sum_{P'\in\mathcal{P}} X^1_{\mathcal{P}}(P') + \frac{k-3}{2}\sum_{P'\in\mathcal{P}} X^2_{\mathcal{P}}(P') + ... + \frac{1}{k-2}\sum_{P'\in\mathcal{P}} X^{k-2}_{\mathcal{P}}(P')}{k-1} \geq$$

$$\geq \frac{k\cdot\ell(G) + k + \frac{1}{k-2}(\sum_{P'\in\mathcal{P}} X^1_{\mathcal{P}}(P') + \sum_{P'\in\mathcal{P}} X^2_{\mathcal{P}}(P') + ... + \sum_{P'\in\mathcal{P}} X^{k-2}_{\mathcal{P}}(P'))}{k-1} \geq$$

$$\geq \frac{k\cdot\ell(G) + k + \frac{k}{k-2}(X^1_{\mathcal{P}}(P) + X^2_{\mathcal{P}}(P) + ... + X^{k-2}_{\mathcal{P}}(P))}{k-1} \geq$$

$$\geq \frac{k(\frac{6f(G,\mathcal{P})}{k-1} + 1) + k + \frac{2k}{k-2}(\frac{2}{k-1}f(G,\mathcal{P}) - 1)}{k-1} = \frac{(6k^2-8k)f(G,\mathcal{P}) + 2k^3 - 8k^2 + 6k}{(k-2)(k-1)^2},$$

and hence $f(G,\mathcal{P}) \leq \frac{(k^3-4k^2+5k-2)n - 2k^3 + 8k^2 - 6k}{6k^2-8k}$. This completes the proof of Theorem 4. $\square$

Proof of Theorem 3. We proceed as in the proof of Theorem 4 and use Corollary 8(i) instead of Lemma 7(i).



Figure 2: Paths S_1, S_2, and S_3

By comparing the lengths of P_2 and S_1 and using Corollary 8(i) and $|V(R)| \geq 2$, we have

$$|V(u_2P_2v)| - 1 \geq |V(Q)| - 1 + |V(R)| - 2 \geq |V(Q)| - 1 \geq f(G, \mathcal{P}).$$

Similarly for S_2 and S_3, we have

$$|V(vP_2x)| - 1 \geq |V(Q)| - 1 + |V(R)| - 1 \geq |V(Q)| \geq f(G, \mathcal{P}) + 1,$$

$$|V(xP_2v_2)| - 1 \geq |V(Q)| - 1 + |V(R)| - 2 \geq |V(Q)| - 1 \geq f(G, \mathcal{P}).$$

Therefore all together we have

$$\ell(G) = |V(P_2)| - 1 = |V(u_2P_2v)| - 1 + |V(vP_2x)| - 1 + |V(xP_2v_2)| - 1 \geq$$

$$\geq f(G, \mathcal{P}) + f(G, \mathcal{P}) + 1 + f(G, \mathcal{P}) = 3f(G, \mathcal{P}) + 1. \qquad (**)$$

By the choice of P and $t'_{\mathcal{P}}(P') \geq 2$ for every $P' \in \mathcal{P}$ together with $(**)$, Lemma 7, and Lemma 6 we have

$$n \geq \frac{4\ell(G) + 4 + 2\sum_{P' \in \mathcal{P}} X^1_{\mathcal{P}}(P') + \frac{1}{2}\sum_{P' \in \mathcal{P}} X^2_{\mathcal{P}}(P')}{3} \geq$$

$$\geq \frac{4(3f(G,\mathcal{P})+1)+4+4(f(G,\mathcal{P})-1)}{3} = \frac{16f(G,\mathcal{P})+4}{3},$$

and hence $f(G,\mathcal{P}) \leq \frac{3n-4}{16}$. This completes the proof of Theorem 3. □

3 Conclusion

As it was mentioned in Introduction, we extend Conjecture 1 to Conjecture 9.

Conjecture 9. *For every connected graph, any k of its longest paths have a common vertex for $3 \leq k \leq 6$.*

Conjecture 10 is an extension of a Conjecture stated in [5] for 3 longest paths. We prove that Conjecture 10 is equivalent with Conjecture 9.

Conjecture 10. *There exists a sublinear function g such that for every connected graph G of order n and every subset $\mathcal{P}$ of $\mathcal{L}(G)$ with $3 \leq |\mathcal{P}| \leq 6$, $f(G,\mathcal{P}) \leq g(n)$.*

Let $\mathcal{G}_n$ be a class of connected graphs of order at least n. In other words, using $d_k(\mathcal{G}_n)$ with $3 \leq k \leq 6$, Conjecture 10 translates into the following statement. The path distance ratio $d_k(\mathcal{G}_n)$ goes to 0 as n goes to infinity.

Theorem 11. *Conjecture 9 is true if and only if Conjecture 10 is true.*

Proof. Suppose Conjecture 9 holds. For every set $\mathcal{P}$ of k longest paths ($3 \leq k \leq 6$) of every connected graph G, we have $f(G,\mathcal{P}) = 0$. Thus any non-negative sublinear function implies that Conjecture 10 holds.

Suppose Conjecture 10 holds. We prove the contrapositive statement, that is, if Conjecture 9 is not true, then neither is Conjecture 10. For $3 \leq k \leq 6$, we consider a connected graph G and a set $\mathcal{P}$ of its k longest paths so that they have no common vertex. We extend G by adding a pendant edge to every vertex, which is an end-vertex of a path of $\mathcal{P}$, and we note that each path of $\mathcal{P}$ prolonged with two of these new edges is a longest path in the extended graph. For a non-negative integer t, we subdivide every edge of the extended graph t times and we observe that the corresponding k paths, say $\mathcal{P}_t$, are longest paths in the resulting graph G_t. Let n be the number of vertices and m the number of edges of G. We see that G_t has at most $n + t(m + 2k)$ vertices. By construction, $f(G_t,\mathcal{P}_t) \geq t$. We consider the sequence of graphs $(G_t)_{t=1}^{\infty}$ and we note that $f(G_t,\mathcal{P}_t)$ cannot be bounded from above by a sublinear function. □

Acknowledgements

This work was partly supported by the project LO1506 of the Czech Ministry of Education, Youth and Sports.

The first and third authors were supported by project GA14-19503S of the Grant Agency of the Czech Republic.

The second author's research is supported by Grant-in-Aid for Scientic Research (C) (15K04979). Also, this work was partially completed during a visit of the second author to the University of West Bohemia. He wishes to express his thanks for the generous hospitality.

References

[1] M. Axenovich, When do 3 longest paths have a common vertex? Discrete Math. Alg. Appl. 1 (2009) 115-120.

[2] P. Balister, E. Győri, J. Lehel, R. Schelp, Longest paths in circular arc graphs, Combin. Probab. Comput. 13 (2004) 311-317.

[3] S. F. de Rezende, C. G. Fernandes, D. M. Martin, Y. Wakabayashi, Intersecting longest paths, Discrete Math. 313 (2013) 1401-1408.

[4] P. Erdös, G. Katona (Eds.), Theory of Graphs, Proceedings of the Colloquium Hald at Tihany, Hungary, 1966, Academic Press, New York, 1968, Problem 4 (T. Gallai), p.362.

[5] S. Fujita, M. Furuya, R. Naserasr, K. Ozeki, A New Approach Towards a Conjecture on Intersecting Three Longest Paths, arXiv:1503.01219v2, 2015.

[6] S. Klavžar, M. Petkovšek, Graphs with nonempty intersection of longest paths, Ars Combin. 26 (1990) 43-52.

[7] Z. Skupień, Smallest sets of longest paths with empty intersection. Combin. Probab. Comput. 5 (1996), no. 4, 429-436.

[8] C. Thomassen, Planar and infinite hypohamiltonian and hypotraceable graphs, Discrete Math. 14 (1976) 377-389.

[9] H. Walther, Über die Nichtexistenz eines Knotenpunktes durch den allen längsten Wege eines Graphen gehen, J. Combin. Theory 6 (1969) 1-6.

[10] H. Walther, H. J. Voss, Über Kreise in Graphen, VED Deutcher Verlag der Wissenschaften, 1974.

[11] T. Zamfirescu, On longest path and circuits in graphs, Math. Scand. 38 (1976) 211-239.

[12] T. Zamfirescu, Intersecting longest path or cycles: short survey, An. Univ. Craiova Ser. Mat. Inform. 28 (2001) 1-9.